

University of Groningen

Decoding therapeutic roles of adipose tissue-derived stromal cells and their extracellular vesicles in liver disease

Afsharzadeh, Danial

DOI:
[10.33612/diss.121499227](https://doi.org/10.33612/diss.121499227)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Afsharzadeh, D. (2020). *Decoding therapeutic roles of adipose tissue-derived stromal cells and their extracellular vesicles in liver disease*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.121499227>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

APPENDIX

A

Summary	142
Nederlandse samenvatting	144
Acknowledgements	148
List of publications	152

SUMMARY

Mesenchymal stromal cells (MSC) are therapeutic cells that, upon the systemic infusion, have the ability to home to the injured liver. MSC improve liver function and ameliorate fibrosis, rendering a promising strategy to treat patients with end-stage liver disease. Although MSC are capable to differentiate into hepatocytes *in vitro*, the MSC-induced therapeutic effects on acute and chronic liver disease appear based on paracrine factors they produce, mediated via the release of trophic factors and extracellular vesicles (EVs). Various trophic molecules have been studied as carriers of the MSC therapeutic effects, while such function for MSC-derived extracellular vesicles has not been yet thoroughly investigated. EVs carry a variety of regulatory nucleic acids and proteins and are able to deliver this cargo to target cells. Cargo of EVs may modulate key cellular processes, such as transcription, post-transcriptional modification and signal transduction in the recipient cells. In this thesis, we established the therapeutic potential of EVs from human adipose tissue-derived stromal cells (hASC-derived EVs) in murine, cellular and tissue (*ex vivo*) models of acute and chronic liver disease.

Chronic liver disease is commonly accompanied by the development of liver fibrosis and hepatic stellate cells (HSC) are considered to be the main drivers of this process that impairs liver function. HSC are located in the space of Disse and are referred as quiescent HSC in the healthy liver. As a result of chronic liver injury, HSC become activated, start to proliferate and produce excessive amounts of extracellular matrix (ECM) proteins, as well as cytokines, chemokines and mitogenic growth factors. Experimental liver fibrosis in rodents is alleviated after systemic administration of MSC yet the type and cellular origin of chemoattractants that cause homing of MSC to the liver are largely unknown. In **Chapter 2**, we found that activated HSC attract hASC to the liver through the secretion of specific chemokines. hASC migrated only to activated HSC and not to quiescent HSC and neither to hepatocytes, highlighting that activation of HSC is a key factor in attraction of hASC to the fibrotic liver. Selective inhibition of chemokine receptors by pharmacological compounds revealed that CXCR2 and CXCR3 are both involved in MSC migration to activated HSC.

hASC are able to suppress liver fibrogenesis, but the underlying mechanisms are largely unknown. In **Chapter 3**, we show that hASC secrete factors (the “hASC-secretome”) that effectively suppress HSC activation *in vitro* and liver fibrosis *in vivo*. These anti-fibrotic properties appear almost exclusively in the EV-fraction, and are not found in the EV-free fraction that contains the trophic factors. These

data indicate that hASC-derived EVs suppress hepatic fibrosis *in vitro*, *in vivo* and *ex vivo*. EVs carry a great variety of different proteins and various types of RNA, such as miRNAs. Our primary miRNA analysis revealed that hASC-derived EVs contain over 1,000 unique types of miRNAs. Pathway enrichment analysis indicated various associations between these miRNAs and pathways regulating fibroblast function and fibrosis, such as MAP Kinase and FoxO signaling pathways.

Acute liver failure is a life-threatening disease where the damaged liver fails within days to weeks. Acute liver damage is associated with inflammation and massive hepatocyte loss. In **Chapter 4**, we challenged the potential of hASC-derived EVs to attenuate acute liver damage in acetaminophen (APAP)- and carbon tetrachloride (CCl₄)-induced murine models of acute liver injury. Both the prophylactic, as well as the therapeutic application of hASC-derived EVs rescued the acute liver injury and suppressed the inflammation in these murine liver disease models. Moreover, hASC-derived EVs suppressed the early onset of liver fibrosis induced by the single high-dose of CCl₄, in line with observations made in Chapter 3.

Hepatic steatosis is the accumulation of fat in the liver. As a consequence of a Western diet, the physiological equilibrium in hepatocyte lipid metabolism is disturbed and uptake and/or *de novo* synthesis of fatty acids is increased relative to fatty acid oxidation. This stimulates the synthesis of triglycerides in the liver to dispose the excess free fatty acids resulting in steatosis. In **Chapter 5**, we show that hASC-derived EVs reverse hepatic steatosis in mice placed on a Western diet for 6 weeks. The 6-week Western diet induced triglyceride accumulation in the liver, but did not cause liver damage, inflammation and fibrosis yet. Weekly administration of hASC-derived EVs in the last 3 weeks of the Western diet feeding lowered hepatic lipid accumulation. Thus, this mild steatosis model of NAFLD showed that hASC-derived EVs directly suppress hepatic steatosis, independent of the co-existence of inflammation and/or fibrosis.

In conclusion, CXCR2 and CXCR3 are prospective targets for the improvement of hASC homing in liver disease. hASC-derived EVs are anti-fibrotic and anti-steatotic, and are able to suppress acute hepatocyte injury. However, mechanisms of these beneficial effects remain to be established in detail in future studies. This thesis provides a platform for the future application of hASC-derived EVs in the treatment of acute and chronic liver disease.

NEDERLANDSE SAMENVATTING

Mesenchymale stromale cellen (MSC) zijn therapeutische cellen die na systemische infusie het vermogen hebben om naar de zieke lever te gaan. MSC verbeteren de leverfunctie en herstellen fibrose, wat een veelbelovende strategie is voor de behandeling van patiënten met vergevorderde leverziekte. Hoewel MSC onder gecontroleerde omstandigheden *in vitro* in levercellen (hepatocyten) kunnen differentiëren, lijken de therapeutische effecten van MSC in acute en chronische leverziekte vooral afkomstig te zijn van paracrine factoren die ze produceren. Effecten van MSC worden gemedieerd via de afgifte van kleine eiwitten (trofische factoren) en extracellulaire vesikels (EV's). De therapeutische effecten van MSC zijn al toegeschreven aan verschillende trofische factoren, maar dergelijke functies zijn nog niet grondig onderzocht voor MSC-geproduceerde EV's. EV's bevatten een verscheidenheid aan regulerende nucleïnezuuren en eiwitten en zijn in staat om deze aan zieke organen af te leveren. De bestanddelen van EV's kunnen belangrijke cellulaire processen moduleren, zoals transcriptie, post-transcriptionele modificatie en signaaltransductie in de ontvangende cellen. In dit proefschrift hebben we het therapeutische potentieel van EV's van MSC onderzocht, in het bijzonder van menselijk vetweefsel afkomstige stromale cellen (hASC-geproduceerde-EV's) in muis-, cel- en weefsel- (*ex vivo*) modellen van acute en chronische leverziekte.

Chronische leverziekte gaat vaak gepaard met de ontwikkeling van leverfibrose. Hepatische stellaatcellen (HSC) worden beschouwd als de belangrijkste oorzaak van dit proces dat de leverfunctie schaadt. HSC bevinden zich in de ruimte van Disse en worden in de gezonde lever aangeduid als "rustende" HSC. Als gevolg van chronisch leverletsel worden HSC geactiveerd en gaan deze cellen prolifereren en produceren ze buitensporige hoeveelheden extracellulaire matrix (ECM) eiwitten, evenals cytokinen, chemokinen en groeifactoren. Systemische toediening van MSC onderdrukt leverfibrose bij knaagdieren, maar het type en de cellulaire oorsprong van chemoattractanten die MSC naar de lever brengen, zijn grotendeels onbekend. In **Hoofdstuk 2** ontdekten we dat geactiveerde HSC specifieke chemokinen uitscheiden die ervoor zorgen dat hASC naar de zieke lever trekken. hASC migreerden alleen naar geactiveerde HSC en niet naar rustende HSC en ook niet naar hepatocyten, wat aangeeft dat activering van HSC een belangrijk proces is voor het aantrekken van hASC naar de fibrotische lever. Selectieve remming van chemokinereceptoren door farmacologische stoffen onthulde dat de receptoren CXCR2 en CXCR3 betrokken zijn bij MSC-migratie naar geactiveerde HSC.

hASC kunnen leverfibrose onderdrukken, maar de onderliggende mechanismen zijn grotendeels onbekend. In **Hoofdstuk 3** laten we zien dat hASC factoren uitscheiden (het 'hASC-secretoom') die HSC-activering *in vitro* en leverfibrose *in vivo* effectief remmen. Deze anti-fibrotische eigenschappen zitten bijna uitsluitend in de EV-fractie en worden niet gevonden in de EV-vrije fractie die de trofische factoren bevat. Deze gegevens geven aan dat hASC- geproduceerde-EV's lever fibrose *in vitro*, *in vivo* en *ex vivo* onderdrukken. EV's bevatten een grote verscheidenheid aan verschillende eiwitten en verschillende soorten RNA, zoals miRNA's. Onze miRNA-analyse onthulde dat hASC- geproduceerde-EV's meer dan 1.000 verschillende soorten miRNA's bevatten. Onze signaleringsroute analyse (pathway enrichment analysis) wees op verschillende associaties tussen deze miRNA's en routes die de fibroblastfunctie en fibrose reguleren, zoals MAP Kinase- en FoxO-signaleringsroutes.

Acuut leverfalen is een levensbedreigende ziekte waarbij de beschadigde lever binnen enkele dagen tot weken faalt. Acute leverschade wordt geassocieerd met ontsteking en massaal verlies van hepatocyten. In **Hoofdstuk 4** hebben we getest of hASC-geproduceerde-EV's ook in staat zijn acute leverschade door paracetamol (APAP) of tetrachloormethaan (CCl_4) in muizen kan tegengaan. Zowel de profylactische als de therapeutische toepassing van hASC-geproduceerde-EV's beschermden de lever tegen de acute leverbeschadiging en onderdrukten de ontsteking in deze muismodellen van leverziekte. Bovendien onderdrukten hASC-geproduceerde-EV's de eerste kenmerken van leverfibrose geïnduceerd door CCl_4 , in overeenstemming met de resultaten beschreven in hoofdstuk 3.

Leververvetting (steatose) is de ophoping van vet in de lever. Als gevolg van een westers dieet wordt het fysiologische evenwicht in het vetmetabolisme van de hepatocyten verstoord en neemt de opname en/of *de novo* synthese van vetzuren toe ten opzichte van de vetzuuroxidatie. Dit stimuleert de vorming van triglyceriden in de lever om de overvloedige vrije vetzuren af te voeren, wat uiteindelijk resulteert in steatose. In **Hoofdstuk 5** laten we zien dat hASC-geproduceerde-EV's leververvetting tegengaan bij muizen die gedurende 6 weken een westers dieet te eten kregen. Het westerse dieet van 6 weken induceerde ophoping van triglyceriden in de lever, maar veroorzaakte nog geen leverschade, ontsteking of fibrose. Wekelijkse toediening van hASC-geproduceerde-EV's in de laatste 3 weken van het westerse dieet verminderde ophoping van lipiden in de lever. Aldus toonde dit milde steatosemodel van niet-alcoholische vette leverziekte

(NAFLD) aan dat hASC-geproduceerde--EV's leversteatose direct onderdrukken, onafhankelijk van de aanwezigheid van ontsteking en/of fibrose.

We kunnen nu dus concluderen dat CXCR2 en CXCR3 factoren zijn die hASC-migratie naar de lever bij leverziekte kunnen stimuleren. Verder zijn hASC-geproduceerde-EV's anti-fibrotisch en anti-steatotisch en kunnen acuut leverletsel onderdrukken. De mechanismen van deze gunstige effecten moeten echter in toekomstige studies verder in detail worden bepaald. Dit proefschrift biedt een platform voor de toekomstige toepassing van hASC-geproduceerde-EV's bij de behandeling van acute en chronische leverziekte.



ACKNOWLEDGMENT

Dear reader, I will cherish this book not only in my future academic career but during my entire life. Therefore, I'm grateful for the time you take browsing through it and reading what I have written; that means a lot to me!

Further, and as you might assume, there are a number of people without whom I would not have been able to complete this research and making it through my PhD degree.

Charismatic, pragmatic and sympathetic is how I describe you, **Klaas Nico**. You have the ability to fascinate a wide range of people; I really admire that. Thanks for believing in me and all the support since the first moment I stepped into your office.

Marco, when I met you for the first time, I knew that I am at the right place! Thanks for keeping on supporting me in every step of this journey, especially during the last few months when finalizing this thesis. I really appreciate your work ethics and learned a great deal from you.

Professor **Ali Canbay**, thank you for being a generous host during my EASL-fellowship period. I appreciate your kindness and the trust you have in me. Your hard working is exemplary!

Lars, working with you and Svenja was more than a simple collaboration. You guys are inspiring! I sincerely enjoy working with you. I'm looking forward to starting new projects together.

Han, you are the most international person I have ever met. Your global academic knowledge is the key to many tough locks. I really admire your positive mindset.

Tjasso, I see your technical tips in different corners of my thesis. I have always enjoyed our chit-chat coffee times, especially our talks about overseas traveling. Thanks for all of them!

Manon, thanks for putting a smile on my face, every time I asked you something. Your soft-natured personality is amazing. Thanks for helping me with logistics and experiments every now and then!

Janette, the passion and determination you bring to the job is admirable! Thanks for always being swift and ready to help!

Professors **Ruud Bank**, **Marco Arese** and **Karen Bieback**, thank you for your time to review my thesis.

Guido K, thanks for always being open to collaborate and answering to my non-ending questions regarding roles of miRNA in fibrosis and tissue-remodeling!

Emma, you are a self-motivated scientist who is pleasant to work with. Thanks for all fun times during our collaboration. Hope you have a good time in France!

Peter, thanks for letting me use facilities of your slice-lab during past years. I enjoyed working with you guys!

Aycha, thanks for all fun chats and organizing those fun beer-evenings!

Henk, and **Trijnie**, thanks for being available and cooperative!

Niels, the fact that we are from different labs was never a barrier whenever your attention was needed. I admire your eagerness to help! Thanks for all assistance and troubleshooting!

Patrick, I still remember how excited I was, when you visualized the EVs as the product of my first isolation, using EM. Thanks for being open and positive to collaborate!

Sebo, **Rutger** and **Ineke**, you did a great job with sequencing the miRNA content of EVs. Thanks for all technical support!

Hans, our corridor chats were fun, I will miss them!

Linda, teaming up to isolate adipose-derived stem cells is how we got to know each other. I admire your skills of team working and organization. Thanks for your consistent support whenever there was a need for stem cells!

Shiva, thanks for your positive vibes even during the most difficult experiments. Wish you the best in your new career!

My old office mates, **Esther** and **Floris**, although our stay in MDL lab had a short overlap, our casual chats were a part of the great experience of my first days at MDL. Wish you both success with your current jobs!

Easy going and happy is how I know you, **Natalia**. Thanks to you and **Cesar** for having me at your cozy Latin nights with your friends!

Turu and **Mengfan**, I have to thank you for your cooperative attitude in our weekly isolation of primary liver cells in the lab. Good job guys!

Iris, **Mark** and **Douwe**, having you guys around in the cell culture lab was always fun. Iris, thanks for those nice ice cream walks!

Sandra and **Fabio**, thanks for sharing your car with Hannah and me on the way from Copenhagen to Groningen! We still talk about your Formula driving, Sandra.

Ali, I appreciate your eagerness for scientific collaborations. Wishing you a successful career in Pakistan!

Anne vE, thanks for all nice days in Groningen. You know how to leave a lasting impression!

Fan and **Chong Zuo**, I appreciate your assistance and sharing your tuned protocol for fatty acid measurement (although I could never read your secret notes in Chinese).

A loud high five to team Vrijmibo, **Onne**, **Maaïke**, **Ana H**, **Rima**, **Lori**, **Hannah W**, **Jan Freak**, **Marleen**, **Joanne**, **Johanna**, **Herson**, **Archi**, **Yana**, **Ying**, **Guido**, **Raphael**, **Dianne**, **Anna B**, **Anna P**, **Emilia**, **Mathilde**, **Hilde**, **Vera**, **Karin**, **Andrea**, **Christy**, **Alfredo**, **Tim** and **Anosh**, for many cozy Friday evenings and creating a positive spirit at work.

Dicky, definitely you are one of the lab mates I hanged out with the most. Especially, our Friday lunches outside of the hospital were quite refreshing. We should keep on organizing those gym times.

Ivo, apparently sitting back to back of each other in the office could not prevent us enjoying casual chats! Certainly, the end of this roommatehood is not the end of this contact.

Dane, you are a great influencer! Thanks for being such an amazing friend during different seasons of life! Let's grow old together!

My dear paranymphs, **Svenja** and **Mirjam**, calling you only colleagues is not true anymore. You both grew to be valuable friends. Svenja, without your efforts, none of the animal studies in this thesis would exist. Thanks for being a great collaborator! I really admire your loyalty and brave personality. Mirjam, getting you to know in the lab was apparently just a start, as we kept on meeting in more fun places such as on the Salsa and Kizomba stage. Thanks for being trustworthy and open-hearted!

My little sister and brother, **Negin** and **David**, thanks for always being supportive and keeping me in the loop of our warm family, even though I'm already away for many years! Let's organize another family-holiday soon!

Dear **Mom** and **dad**, without you, I would not stand here today. There are no words to thank you enough for your consistent and unconditional support.

Dear **Hannah**, I appreciate your understanding and companionship for many weekends that I sacrificed to work during the past couple of months. Although most of my PhD period had been already passed before we met, I can see your ideas and remarks in almost all corners of my thesis. Thanks for being you!

Dear **Oscar**, my hasty son! Your name was not supposed to be included in this acknowledgment, but by coming earlier than expected, you made it in here, congratulations! The color of light you have shed on our life is wonderful! We love you!

LIST OF PUBLICATIONS

Afsharzadeh D, Sydor S, Gore E, Friedrich J, Krenning G, Van Rijn P, Olinga P, Canbay A, Bechmann LP, Harmsen MC and Faber KN. Adipose tissue-derived stromal cells suppress liver fibrosis in an extracellular vesicle-mediated fashion. *Submitted*

Afsharzadeh D, Sydor S, Saeed A, Canbay A, Olinga P, Bechmann LP, Harmsen MC and Faber KN. Adipose tissue-derived stromal cells are attracted to activated hepatic stellate cells through CXCR2- and CXCR3-mediated signalling. *Submitted*

Afsharzadeh D, Sydor S, Canbay A, Bechman LP, Harmsen MC and Faber KN. Extracellular vesicles from adipose tissue-derived stromal cells ameliorate APAP- and CCl4-induced acute liver injury. *In preparation*

Afsharzadeh D, Sydor S, Canbay A, Bechman LP, Harmsen MC and Faber KN. Extracellular vesicles from adipose tissue-derived stromal cells ameliorate western diet-induced NAFLD in mice. *In preparation*

Nouri HR, Sankian M, **Afsharzadeh D**, Varasteh A. Immunotherapy with a recombinant hybrid molecule alleviates allergic responses more efficiently than an allergenic cocktail or pollen extract in a model of *Chenopodium album* allergy. *Int Arch Allergy Immunol*. 2013.

Tayari M, **Afsharzadeh D**. Amplification of antioxidant activity of haptoglobin (2-2)-hemoglobin at pathologic temperature and presence of antibiotics. *Indian J Clin Biochem*. 2012.

Keshel SH, Soleimani M, Tavirani MR, Ebrahimi M, Raeisossadati R, Yasaei H, **Afsharzadeh D**, Behroz MJ, Atashi A, Amanpour S, Khoshzaban A, Roozafzoon R, Behrouzi GR. Evaluation of unrestricted somatic stem cells as a feeder layer to support undifferentiated embryonic stem cells. *Mol Reprod Dev*. 2012.

Nouri HR, Varasteh A, Vahedi F, Chamani J, **Afsharzadeh D**, Sankian M. Constructing a hybrid molecule with low capacity of IgE binding from *Chenopodium album* pollen allergens. *Immunol Lett*. 2012.

Nouri HR, Sankian M, Vahedi F, **Afsharzadeh D**, Rouzbeh L, Moghadam M, Varasteh A. Diagnosis of *Chenopodium album* allergy with a cocktail of recombinant allergens as a tool for component-resolved diagnosis. *Mol Biol Rep*. 2012.